

RINVOQ is a JAK inhibitor for adults with active psoriatic arthritis (PsA) in whom TNF blockers did not work well.



If you're still experiencing **psoriatic arthritis** symptoms after a TNF blocker like **HUMIRA®** or **ENBREL®**,

RINVOQ IS
DIFFERENT
AND MAY HELP

WHAT IS RINVOQ?

RINVOQ is a prescription medicine used to treat adults with active psoriatic arthritis when 1 or more medicines called tumor necrosis factor (TNF) blockers have been used, and did not work well or could not be tolerated. It is not known if RINVOQ is safe and effective in children with psoriatic arthritis.

SAFETY CONSIDERATIONS

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower ability to fight infections. Serious infections, some fatal, occurred, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses.
- **Increased risk of death in people age 50+ with at least 1 heart disease risk factor.**
- **Cancer and immune system problems.** Increased risk of some cancers, including lymphoma and skin. Current or past smokers have higher risk for lymphoma and lung cancer.
- **Increased risk of major cardiovascular events such as heart attack, stroke, or death in people 50+ with at least 1 heart disease risk factor, especially in current or past smokers.**
- **Blood clots,** some fatal, in veins of the legs or lungs and arteries. This occurred more often in people 50+ with at least 1 heart disease risk factor.
- **Serious allergic reactions. Do not take if allergic to RINVOQ or its ingredients.**
- **Tears in the stomach or intestines; changes in certain laboratory test results.**

Please see full Important Safety Information on pages 8–9. Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

PROVEN

IN CLINICAL STUDIES FOR PsA


RINVOQ can:

- > **Significantly relieve joint pain, swelling, and stiffness**
- > **Stop further irreversible joint damage**
- > **Significantly reduce PsA fatigue**
- > **Leave skin noticeably clearer for people with PsA skin plaques***



Illustration of skin clearance for people with moderate to severe PsA skin plaques. Individual results may vary.

*RINVOQ has not been studied in and is not indicated for treating plaque psoriasis.



- > **Make it easier to do everyday things**
- > **Help people achieve Minimal Disease Activity— that can mean less joint pain, swelling, stiffness, and skin symptoms**
- > **Relieve tenderness for people with enthesitis (pain and swelling in areas like the heel and elbow)**
- > **Reduce swelling of fingers and toes for people with dactylitis (also called “sausage fingers”)**

TALK TO YOUR RHEUMATOLOGIST ABOUT
RINVOQ RELIEF

SAFETY CONSIDERATIONS

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower ability to fight infections. Serious infections, some fatal, occurred, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses.
- **Increased risk of death in people age 50+ with at least 1 heart disease risk factor.**
- **Cancer and immune system problems.** Increased risk of some cancers, including lymphoma and skin. Current or past smokers have higher risk for lymphoma and lung cancer.
- **Increased risk of major cardiovascular events such as heart attack, stroke, or death in people 50+ with at least 1 heart disease risk factor, especially in current or past smokers.**
- **Blood clots,** some fatal, in veins of the legs or lungs and arteries. This occurred more often in people 50+ with at least 1 heart disease risk factor.
- **Serious allergic reactions. Do not take if allergic to RINVOQ or its ingredients.**
- **Tears in the stomach or intestines; changes in certain laboratory test results.**

Please see full Important Safety Information on pages 8–9. Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

RINVOQ HELPS RELIEVE

PHYSICAL AND MENTAL PsA SYMPTOMS

In clinical studies, RINVOQ helped improve both physical and mental PsA symptoms at 12 weeks.

That's according to a survey called the **36-Item Short Form Health Survey (SF-36)**, which asked people with PsA 36 questions about their health and quality of life while taking RINVOQ.



WHAT DOES THE SF-36 SURVEY ASK ABOUT?



PHYSICAL TOPICS

- **Everyday impact of PsA**
Time and effort to do daily things
- **Pain**
Both its intensity and its everyday toll
- **Physical function**
Ability to do things like climb stairs, get dressed, or do housework
- **General health**
And overall wellness



MENTAL TOPICS

- **Vitality**
Time spent feeling energetic vs tired
- **Social functioning**
Taking part in social activities
- **Emotional impact**
Time and care for work and everyday tasks
- **Mental health**
Time spent feeling happy, calm, nervous, or sad

SAFETY CONSIDERATIONS

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower ability to fight infections. Serious infections, some fatal, occurred, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses.
- **Increased risk of death in people age 50+ with at least 1 heart disease risk factor.**
- **Cancer and immune system problems.** Increased risk of some cancers, including lymphoma and skin. Current or past smokers have higher risk for lymphoma and lung cancer.
- **Increased risk of major cardiovascular events such as heart attack, stroke, or death in people 50+ with at least 1 heart disease risk factor, especially in current or past smokers.**
- **Blood clots,** some fatal, in veins of the legs or lungs and arteries. This occurred more often in people 50+ with at least 1 heart disease risk factor.
- **Serious allergic reactions. Do not take if allergic to RINVOQ or its ingredients.**
- **Tears in the stomach or intestines; changes in certain laboratory test results.**

Please see full Important Safety Information on pages 8–9. Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

MORE RINVOQ RESULTS

FAST RELIEF THAT CAN LAST

In a clinical study, RINVOQ relieved PsA joint pain, swelling, and stiffness for some people **in as little as 2 weeks**.

The majority of people taking RINVOQ had significantly improved symptoms **at 12 weeks**.

A similar percentage of people reported relief **over a year later** (at 56 weeks).*

Helps relieve PsA joint pain, swelling, & stiffness

2 weeks | **12** weeks | **56** weeks

Individual results may vary.



RELIEF AT 3 YEARS (152 WEEKS*)

Of those who continued on RINVOQ for 3 years, many saw improvement in PsA symptoms.

In this less rigorous part of the study, patients and physicians were aware of RINVOQ treatment, which may have influenced the results at 3 years.

*20% improvement in PsA symptoms when used with or without csDMARDs. Doctors adjusted supportive PsA medicines as needed at week 16 or 36.

SAFETY CONSIDERATIONS

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower ability to fight infections. Serious infections, some fatal, occurred, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses.
- **Increased risk of death in people age 50+ with at least 1 heart disease risk factor.**
- **Cancer and immune system problems.** Increased risk of some cancers, including lymphoma and skin. Current or past smokers have higher risk for lymphoma and lung cancer.
- **Increased risk of major cardiovascular events such as heart attack, stroke, or death in people 50+ with at least 1 heart disease risk factor, especially in current or past smokers.**
- **Blood clots,** some fatal, in veins of the legs or lungs and arteries. This occurred more often in people 50+ with at least 1 heart disease risk factor.
- **Serious allergic reactions. Do not take if allergic to RINVOQ or its ingredients.**
- **Tears in the stomach or intestines; changes in certain laboratory test results.**

Please see full Important Safety Information on pages 8–9. Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

A ONCE-DAILY PILL NOT AN INJECTION

- > Take RINVOQ once a day, every day
- > You can take RINVOQ with or without food
 - Avoid food or drink containing grapefruit
 - Do not split, crush, or chew the pill
- > Your doctor may prescribe RINVOQ with or without a nonbiologic disease-modifying antirheumatic drug (DMARD) like methotrexate
- > You and your doctor will determine your optimal treatment plan

Call your healthcare provider if you see medication residue (a whole tablet or tablet pieces) in your stool.

EASY-TO-OPEN BOTTLE

Awarded the Arthritis Foundation's Ease of Use Commendation, RINVOQ's innovative bottle cap includes an EASY-OPEN GRIP and embedded tool to puncture the foil liner.

See how the bottle opens at RINVOQ.com/psa-bottle



RINVOQ in ACTION

With psoriatic arthritis, your **overactive immune system** attacks your joints and skin.*

*RINVOQ is not approved to treat plaque psoriasis.



RINVOQ is a JAK inhibitor that **REGULATES** your overactive immune system to **HELP STOP** the attack.

RINVOQ works inside your cells to block certain signals that are thought to cause inflammation.

Learn more about the science behind RINVOQ and how it works at RINVOQ.com/psa-science

SAFETY CONSIDERATIONS

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower ability to fight infections. Serious infections, some fatal, occurred, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses.
- **Increased risk of death in people age 50+ with at least 1 heart disease risk factor.**
- **Cancer and immune system problems.** Increased risk of some cancers, including lymphoma and skin. Current or past smokers have higher risk for lymphoma and lung cancer.
- **Increased risk of major cardiovascular events such as heart attack, stroke, or death in people 50+ with at least 1 heart disease risk factor, especially in current or past smokers.**
- **Blood clots,** some fatal, in veins of the legs or lungs and arteries. This occurred more often in people 50+ with at least 1 heart disease risk factor.
- **Serious allergic reactions. Do not take if allergic to RINVOQ or its ingredients.**
- **Tears in the stomach or intestines; changes in certain laboratory test results.**

Please see full Important Safety Information on pages 8–9. Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.



For adults with active psoriatic arthritis in whom TNF blockers did not work well.

HOW TO TALK TO YOUR RHEUM

PsA can be an unpredictable condition, and your response to treatment may change over time. So it's important that your rheumatologist knows how you're doing. Not just your goals and progress, but the setbacks and frustrations, too. All the ways your PsA symptoms make it hard to get back to the things that matter.

Be honest • Be specific • Be candid

No detail is too big or small. Tell your rheum how you really feel, and make sure that they understand what you tell them. That way they'll have a complete picture of your life with PsA, so you can decide on your best path forward, together.

Remember, results don't happen overnight. But when you have a doctor who listens, you have the best chance to reach your goals.



> **Get ready for your next talk with our appointment checklist**

Visit [RINVOQ.com/psa-checklist](https://rinvoq.com/psa-checklist)

> **Find a rheumatologist near you**

Visit [RINVOQ.com/psa-rheum](https://rinvoq.com/psa-rheum)

TAKE ON PSA

ASK YOUR RHEUMATOLOGIST FOR RINVOQ

RINVOQ is not for everyone. Before you and your doctor make a treatment decision, read about the Use and Important Safety Information on pages 8–9. Working together with your doctor can help you determine if RINVOQ is right for you.

Refer to full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

WE'RE IN YOUR CORNER



Once you and your doctor decide to take on PsA with **RINVOQ**, you'll have access to **RINVOQ Complete**—a one-to-one personalized experience that provides the resources you need, when you need them.


Regardless of your insurance coverage or status, your **RINVOQ Complete Nurse Ambassador*** can connect you with an Insurance Specialist who can help you find ways to save on your medication.


>95% of people with self- or employer-provided insurance plans or Medicare Part D are covered for RINVOQ†


You may pay as little as **\$5/mo*** with the RINVOQ Complete Savings Card, if eligible

RINVOQ[®] COMPLETE

To enroll you can:

 Work with your **doctor's office**

 Text "ENROLL" or call **1.800.2RINVOQ** (1.800.274.6867)
1 msg/request. Msg & Data Rates May Apply. Visit www.abbvie.com/privacy.html for info on AbbVie's privacy practices, and SeeRTerms.com for terms and conditions.

 Visit **RINVOQComplete.com** for personalized support or to chat live with a RINVOQ Complete agent

Trouble affording your medication?
 See if AbbVie can help.
 Call us at **1.800.2RINVOQ**

*Nurse Ambassadors are provided by AbbVie and do not work under the direction of your health care professional (HCP) or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

†National commercial and Medicare Part D formulary coverage as of May 2023.

Eligibility: Available to patients with commercial insurance coverage for RINVOQ[®] (upadacitinib) who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. **For full Terms and Conditions**, visit RinvoqSavingsCard.com or call **1.800.2RINVOQ** for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit <https://privacy.abbvie>

Please see Important Safety Information on pages 8–9. Refer to full **Prescribing Information**, including **Medication Guide**, at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.



For adults with active psoriatic arthritis in whom TNF blockers did not work well.

USE AND IMPORTANT SAFETY INFORMATION

WHAT IS RINVOQ?

RINVOQ is a prescription medicine used to treat adults with active psoriatic arthritis when 1 or more medicines called tumor necrosis factor (TNF) blockers have been used, and did not work well or could not be tolerated. It is not known if RINVOQ is safe and effective in children with psoriatic arthritis.

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT RINVOQ?

RINVOQ may cause serious side effects, including:

- **Serious infections.** RINVOQ can lower your ability to fight infections. Serious infections have happened while taking RINVOQ, including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections. Your healthcare provider (HCP) should test you for TB before starting RINVOQ and check you closely for signs and symptoms of TB during treatment with RINVOQ. You should not start taking RINVOQ if you have any kind of infection unless your HCP tells you it is okay. If you get a serious infection, your HCP may stop your treatment until your infection is controlled. You may be at higher risk of developing shingles (herpes zoster).
- **Increased risk of death in people 50 years and older who have at least 1 heart disease (cardiovascular) risk factor.**
- **Cancer and immune system problems.** RINVOQ may increase your risk of certain cancers. Lymphoma and other cancers, including skin cancers, can happen. Current or past smokers are at higher risk of certain cancers, including lymphoma and lung cancer. Follow your HCP's advice about having your skin checked for skin cancer during treatment with RINVOQ. Limit the amount of time you spend in sunlight. Wear protective clothing when you are in the sun and use sunscreen.
- **Increased risk of major cardiovascular (CV) events, such as heart attack, stroke, or death, in people 50 years and older who have at least 1 heart disease (CV) risk factor, especially if you are a current or past smoker.**
- **Blood clots.** Blood clots in the veins of the legs or lungs and arteries can happen with RINVOQ. This may be life-threatening and cause death. Blood clots in the veins of the legs and lungs have happened more often in people who are 50 years and older and with at least 1 heart disease (CV) risk factor.
- **Allergic reactions.** Symptoms such as rash (hives), trouble breathing, feeling faint or dizzy, or swelling of your lips, tongue, or throat, that may mean you are having an allergic reaction have been seen in people taking RINVOQ. Some of these reactions were serious. If any of these symptoms occur during treatment with RINVOQ, stop taking RINVOQ and get emergency medical help right away.
- **Tears in the stomach or intestines.** This happens most often in people who take nonsteroidal anti-inflammatory drugs (NSAIDs) or corticosteroids. Get medical help right away if you get stomach-area pain, fever, chills, nausea, or vomiting.
- **Changes in certain laboratory tests.** Your HCP should do blood tests before you start taking RINVOQ and while you take it. Your HCP may stop your RINVOQ treatment for a period of time if needed because of changes in these blood test results.

Do not take RINVOQ if you are allergic to upadacitinib or any of the ingredients in RINVOQ. See the Medication Guide or Consumer Brief Summary for a complete list of ingredients.

WHAT SHOULD I TELL MY HCP BEFORE STARTING RINVOQ?

Tell your HCP if you:

- Are being treated for an infection, have an infection that won't go away or keeps coming back, or have symptoms of an infection, such as:
 - Fever, sweating, or chills
 - Shortness of breath
 - Warm, red, or painful skin or sores on your body
 - Muscle aches
 - Feeling tired
 - Blood in phlegm
 - Diarrhea or stomach pain
 - Cough
 - Weight loss
 - Burning when urinating or urinating more often than normal
- Have TB or have been in close contact with someone with TB.
- Are a current or past smoker.
- Have had a heart attack, other heart problems, or stroke.
- Have or have had any type of cancer, hepatitis B or C, shingles (herpes zoster), blood clots in the veins of your legs or lungs, diverticulitis (inflammation in parts of the large intestine), or ulcers in your stomach or intestines.
- Have other medical conditions, including liver problems, low blood cell counts, diabetes, chronic lung disease, HIV, or a weak immune system.
- Live, have lived, or have traveled to parts of the country, such as the Ohio and Mississippi River valleys and the Southwest, that increase your risk of getting certain kinds of fungal infections. If you are unsure if you've been to these types of areas, ask your HCP.
- Have recently received or are scheduled to receive a vaccine. People who take RINVOQ should not receive live vaccines.
- Are pregnant or plan to become pregnant. Based on animal studies, RINVOQ may harm your unborn baby. Your HCP will check whether or not you are pregnant before you start RINVOQ. You should use effective birth control (contraception) to avoid becoming pregnant during treatment with RINVOQ and for 4 weeks after your last dose.
- There is a pregnancy surveillance program for RINVOQ. The purpose of the program is to collect information about the health of you and your baby. If you become pregnant while taking RINVOQ, you are encouraged to report the pregnancy by calling 1-800-633-9110.
- Are breastfeeding or plan to breastfeed. RINVOQ may pass into your breast milk. Do not breastfeed during treatment with RINVOQ and for 6 days after your last dose.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RINVOQ and other medicines may affect each other, causing side effects.

Especially tell your HCP if you take:

- Medicines for fungal or bacterial infections
- Rifampicin or phenytoin
- Medicines that affect your immune system

If you are not sure if you are taking any of these medicines, ask your HCP or pharmacist.

CONTINUED ON NEXT PAGE

IMPORTANT SAFETY INFORMATION

(continued from previous page)

WHAT SHOULD I AVOID WHILE TAKING RINVOQ?

Avoid food or drink containing grapefruit during treatment with RINVOQ as it may increase the risk of side effects.

WHAT SHOULD I DO OR TELL MY HCP AFTER STARTING RINVOQ?

- Tell your HCP right away if you have any symptoms of an infection. RINVOQ can make you more likely to get infections or make any infections you have worse.
- Get emergency help right away if you have any symptoms of a heart attack or stroke while taking RINVOQ, including:
 - Discomfort in the center of your chest that lasts for more than a few minutes or that goes away and comes back
 - Severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw
 - Pain or discomfort in your arms, back, neck, jaw, or stomach
 - Shortness of breath with or without chest discomfort
 - Breaking out in a cold sweat
 - Nausea or vomiting
 - Feeling lightheaded
 - Weakness in one part or on one side of your body
 - Slurred speech
- Tell your HCP right away if you have any signs or symptoms of blood clots during treatment with RINVOQ, including:
 - Swelling
 - Pain or tenderness in one or both legs
 - Sudden unexplained chest or upper back pain
 - Shortness of breath or difficulty breathing
- Tell your HCP right away if you have a fever or stomach-area pain that does not go away, and a change in your bowel habits.

WHAT ARE OTHER POSSIBLE SIDE EFFECTS OF RINVOQ?

Common side effects include upper respiratory tract infections (common cold, sinus infections), shingles (herpes zoster), herpes simplex virus infections (including cold sores), bronchitis, nausea, cough, fever, acne, headache, increased blood levels of creatine phosphokinase, allergic reactions, inflammation of hair follicles, stomach-area (abdominal) pain, increased weight, flu, tiredness, lower number of certain types of white blood cells (neutropenia, lymphopenia, leukopenia), muscle pain, flu-like illness, rash, increased blood cholesterol levels, increased liver enzyme levels, pneumonia, low number of red blood cells (anemia), and infection of the stomach and intestine (gastroenteritis).

A separation or tear to the lining of the back part of the eye (retinal detachment) has happened in people with atopic dermatitis treated with RINVOQ. Call your HCP right away if you have any sudden changes in your vision during treatment with RINVOQ.

Some people taking RINVOQ may see medicine residue (a whole tablet or tablet pieces) in their stool. If this happens, call your healthcare provider.

These are not all the possible side effects of RINVOQ.

HOW SHOULD I TAKE RINVOQ?

RINVOQ is taken once a day with or without food. Do not split, crush, or chew the tablet. Take RINVOQ exactly as your HCP tells you to use it. RINVOQ is available in 15 mg, 30 mg, and 45 mg extended-release tablets.

This is the most important information to know about RINVOQ. For more information, talk to your HCP.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Please see full [Prescribing Information](#), including [Medication Guide](#), at www.rxabbvie.com/pdf/rinvoq_pi.pdf and discuss with your doctor.

© 2023 AbbVie. All rights reserved.
US-RNQR-230122 June 2023



ASK YOUR RHEUMATOLOGIST FOR
RINVOQ

**SIGN UP FOR ALL THE LATEST
ABOUT RINVOQ**